

公司产品仅供科学研究实验, 不得用于临床!

商品详情:

英文名称: ox-LDL

中文名称: **氧化低密度脂蛋白单克隆抗体**

别名: ox-LDL; LDL (Copper oxidized); Cu₂SO₄ oxidized low density lipoprotein; oxidized low density lipoprotein; Low density lipoprotein; MDA oxidized LDL; MDA oxidized low density lipoprotein.

研究领域: 心血管 细胞生物 免疫学 脂蛋白

抗体来源: Mouse

克隆类型: Monoclonal

克隆号: 5C1

交叉反应: Mouse, (predicted: Human,)

产品应用: WB=1:500-2000 ELISA=1:5000-10000 IHC-P=1:100-500 IHC-F=1:100-500 ICC=1:100-500 IF=1:100-500 (石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

理论分子量: 31kDa

细胞定位: 细胞膜 分泌型蛋白

性状: Liquid

浓度: 1mg/ml

免疫原: Full length protein from human plasma

亚型: IgG

缓冲液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件: Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.

注意事项: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

氧化低密度脂蛋白单克隆抗体产品介绍: Low-density lipoprotein (LDL) is the carrier protein for cholesterol in the blood. LDL binds to its receptor on the capillary walls and thereby mediates the uptake and clearance of cholesterol from the circulation. In atherosclerotic lesions oxidatively modified LDL is found and oxidized LDL is specifically recognized and ingested by macrophages via scavenger receptor A and CD36. Oxidized LDL may be a marker of atherosclerosis but the precise changes in oxidized LDL are not well described. Low-density lipoprotein oxidised with Cu₂SO₄.

When too much LDL cholesterol circulates in the blood, it can slowly build up in the inner walls of the arteries that feed the heart and brain. Together with other substances it can form plaque, a thick, hard deposit that can clog those arteries. This

condition is known as atherosclerosis. Oxidized lipoproteins are formed by free radical damage to lipids that accumulate in macrophages and smooth muscle cells causing foam cell formation, an initial step in the disease.

www.pyram.cn